



Optimizing quality and shelf-life of retail pork cuts by using muscle specific three-gas MA-packaging

Mari Ann Tørngren, Mianne Darré and Lars Kristensen
Danish Meat Research Institute, DMRI, Roskilde, Denmark

INTRODUCTION

Packaging of fresh pork in high oxygen MAP can have negative effects on meat quality. This results in less tender and less juicy meat with a rancid off-flavor compare to wrapped or vacuum packed meat. The objective of this study was to investigate the effect of low oxygen three-gas MAP on shelf-life and eating quality.

MATERIALS AND METHOD

Retail cuts from *M. longissimus dorsi* (LD), *M. semimembranosus* (SM), and pork belly were packed in five different gas compositions, stored at 5°C and analyzed during storage until day 13.

For each cut, the experimental work was divided into two sub-trials – part 1: meat quality and part 2: shelf-life. For both sub-trials, all pigs were slaughtered on the same date, but for Part 1 pigs were selected according to gender (female) and weight (79-83 kg), whereas random pigs were used for Part 2.

The same slaughter process was used for all three cuts: slaughtering on Monday, and cutting, deboning, shell freezing on Tuesday. For part 1 the meat was sliced at DMRI on Wednesday just before packaging, whereas meat for Part 2 was sliced on Tuesday at the slaughterhouse. All meat samples were packed in modified atmosphere at DMRI on Wednesday in the following five gas compositions.

GAS	O ₂	CO ₂	N ₂
0/20	0%	20%	80%
40/20	40%	20%	40%
50/20	50%	20%	30%
50/40	50%	40%	10%
80/20	80%	20%	0%

RESULTS

Only results for chops and belly are shown.

Table 1. Sensory attributes of pan-cooked pork chops (LD) in MAP with different gas mixtures (day 9, O₂/CO₂).

	0/20	40/20	50/20	50/40	80/20
APPEARANCE (day 9)					
SHELF-LIFE - COLOUR	>13 days	>13 days	>13 days	9-12 days	9-12 days
SHELF-LIFE - ODOUR	9-12 days	9-12 days	9-12 days	9-12 days	13 days
PMB (day 2)					
DONENESS	6.2 c	7.9 b	7.8 b	8.6 a	8.7 a
TENDERNESS	6.1 a	5.6 ab	5.3 bc	4.7 c	5.6 ab
JUICINESS	6.1 a	5.9 a	6.1 a	4.9 b	5.6 a
WOF	1.3 c	2.0 bc	1.4 c	2.4 ab	2.8 a

Table 2. Sensory attributes of pan-cooked pork belly in MAP with different gas mixtures (day 7, O₂/CO₂).

	0/20	40/20	50/20	50/40	80/20
APPEARANCE (Day 7)					
SHELF-LIFE - COLOUR	> 9 days	> 9 days	> 9 days	> 9 days	> 9 days
SHELF-LIFE - ODOUR	5-6 days	3-4 days	5-6 days	5 days	5-6 days
CRISPNESS	4.9 b	6.5 a	6.4 a	7.1 a	3.2 c
RANCID	1.0	1.3	0.9	1.2	1.6
STALE FLAVOUR	4.0 a	4.0 a	3.2 b	2.7 b	4.6 a
BITTER TASTE	5.0 a	4.8 ab	4.4 b	4.7 ab	4.6 ab

CONCLUSION

Gas compositions in modified atmosphere packaging of retail packed pork cuts must be muscle-specific in order to optimise shelf-life, colour stability and eating quality.

Packaging of pork chops and schnitzels in 40% O₂ + 20% CO₂ + 40% N₂ maintained the same shelf-life as traditional MAP 80% O₂ + 20% CO₂ and resulted in more tender and juicy meat with less PMB and rancid flavour. Packaging of sliced pork belly in 50% O₂ + 40% CO₂ + 10% N₂ resulted in a more crispy texture and less stale and bitter taste. Three-gas MAP could therefore be a suitable alternative to traditional MAP, maintaining shelf-life and enhancing eating quality.



CONTACT

LARS KRISTENSEN | LRK@DTI.DK | +45 7220 2670